

Key to species of Nearctic *Brachycyrtus* (Hymenoptera: Ichneumonidae: Brachycyrtinae)

by David B. Wahl
[adapted from Townes (1960)]

1. Supraclypeal area completely white; eyes with deep impressions opposite antennal sockets (Fig. 1). Hind leg as in Fig. 2. Propodeum with areola separated from petiolar area by carina (Fig. 03). Setae of metapleuron and propodeum longer than mesoscutal setae (Fig. 04).
..... ***pretiosus*** Cushman
- 1'. Supraclypeal area with varying amounts of dark brown/fuscous and yellow; eyes with impressions opposite antennal sockets more shallow (Figs. 5-8). Hind leg as in Fig. 9. Propodeum with areola confluent with petiolar area (Fig. 10). Setae of metapleuron and propodeum about same length as mesoscutal setae (Fig. 11).
..... ***ornatus*** Kriechbaumer

Figures

The American Entomological Institute photograph voucher code for an individual specimen follows the species name. For example, '#0103-04' is the fourth photograph taken of voucher specimen 103. All specimens are in the American Entomological Institute collection unless otherwise noted.

- Fig. 01 - *Brachycyrtus pretiosus* #0103-04
- Fig. 02 - *Brachycyrtus pretiosus* #0093-02
- Fig. 03 - *Brachycyrtus pretiosus* #0094-01
- Fig. 04 - *Brachycyrtus pretiosus* #0103-03
- Fig. 05 - *Brachycyrtus ornatus* #0107-01
- Fig. 06 - *Brachycyrtus ornatus* #0097-04
- Fig. 07 - *Brachycyrtus ornatus* #0096-03
- Fig. 08 - *Brachycyrtus ornatus* #0106-02
- Fig. 09 - *Brachycyrtus ornatus* #0097-01
- Fig. 10 - *Brachycyrtus ornatus* #0098-01
- Fig. 11 - *Brachycyrtus ornatus* #0102-02

Key and images © American Entomological Institute, following Attribution-NonCommercial 2.5 license (CC BY-NC 2.5) ([link](http://creativecommons.org/licenses/by-nc/2.5/): <http://creativecommons.org/licenses/by-nc/2.5/>). Please cite this work as: Genera Ichneumonorum Nearctiae. 2014 (and updates). Version: 2014.09.04. <http://www.amentinst.org/GIN/>.