

Key to species of New World *Stilbops* (Hymenoptera: Ichneumonidae: Stilbopinae)

by David B. Wahl
[(adapted from Townes (1989)]

1. Ventral surface of flagellum yellowish-brown or light brown (Fig. 1). T5-9 dark brown to fuscous in both sexes.
T2-3 of female mostly to entirely brownish-red (Fig. 2), black in male (Fig. 3). Female with propodeal carinae complete and strong (Fig. 4). Distribution: northeastern North America.
..... ***latibasis*** Townes
- 1'. Flagellum uniformly dark brown (Fig. 5). T6-9 brownish-red in female, black in male. T2-3 black in both sexes (Figs. 6-7). Female with propodeal carina weak; posterior transverse carina absent and longitudinal carinae apically absent (Fig. 8). Distribution: southern Mexico (Oaxaca).
..... ***mexicana*** Townes

Figures

The American Entomological Institute photograph voucher code for an individual specimen follows the species name. For example, '#0146-03' is the third photograph taken of voucher specimen 146. All specimens are in the American Entomological Institute collection unless otherwise noted.

- Fig. 01 - *Stilbops latibasis* 0146-03
- Fig. 02 - *Stilbops latibasis* 0146-04
- Fig. 03 - *Stilbops latibasis* [Canadian National Collection of Insects, Ottawa, Canada]
- Fig. 04 - *Stilbops latibasis* 0146-06
- Fig. 05 - *Stilbops mexicana* 0147-03
- Fig. 06 - *Stilbops mexicana* 0147-02
- Fig. 07 - *Stilbops mexicana* 0148-01
- Fig. 08 - *Stilbops mexicana* 0147-04

Key and images © American Entomological Institute, following Attribution-NonCommercial 2.5 license (CC BY-NC 2.5) (link: <http://creativecommons.org/licenses/by-nc/2.5/>). Please cite this work as: Genera Ichneumonorum Nearcticae. 2014 (and updates). Version: 2014.09.04. <http://www.amentinst.org/GIN/>.